

Hidden symmetries of the Nambu-Goto action

M. J. Duff¹

*The Blackett Laboratory, Imperial College London,
Prince Consort Road, London SW7 2AZ*

ABSTRACT

We organize the eight variables of the four-dimensional bosonic string $(\dot{X}^\mu, X'^\mu)$ into a $2 \times 2 \times 2$ hypermatrix $a_{AA'A''}$ and show that in signature $(2, 2)$ the Nambu-Goto Lagrangian is given by $\sqrt{\text{Det}} a$ where Det is Cayley's hyperdeterminant. This is invariant not only under $[SL(2, R)]^3$ but also under interchange of the indices A , A' and A'' . This triality reveals hitherto hidden discrete symmetries of the Nambu-Goto action.

¹m.duff@imperial.ac.uk

1 Cayley's hyperdeterminant

In 1845 Cayley [1] generalized the determinant of a 2×2 matrix $a_{AA'}$

$$\begin{aligned}\det a &= \frac{1}{2} \epsilon^{AB} \epsilon^{A'B'} a_{AA'} a_{BB'} \\ &= a_{00} a_{11} - a_{01} a_{10}\end{aligned}\tag{1.1}$$

to the *hyperdeterminant* of a $2 \times 2 \times 2$ *hypermatrix* $a_{AA'A''}$

$$\begin{aligned}\text{Det } a &= -\frac{1}{2} \epsilon^{AB} \epsilon^{A'B'} \epsilon^{CD} \epsilon^{C'D'} \epsilon^{A''D''} \epsilon^{B''C''} a_{AA'A''} a_{BB'B''} a_{CC'C''} a_{DD'D''} \\ &= a_{000}^2 a_{111}^2 + a_{001}^2 a_{110}^2 + a_{010}^2 a_{101}^2 + a_{100}^2 a_{011}^2 \\ &\quad - 2(a_{000} a_{001} a_{110} a_{111} + a_{000} a_{010} a_{101} a_{111} \\ &\quad + a_{000} a_{100} a_{011} a_{111} + a_{001} a_{010} a_{101} a_{110} \\ &\quad + a_{001} a_{100} a_{011} a_{110} + a_{010} a_{100} a_{011} a_{101}) \\ &\quad + 4(a_{000} a_{011} a_{101} a_{110} + a_{001} a_{010} a_{100} a_{111})\end{aligned}\tag{1.2}$$

The hyperdeterminant vanishes iff the following system of equations in six unknowns $p^A, q^{A'}, r^{A''}$ has a nontrivial solution, not allowing any of the pairs to be both zero:

$$\begin{aligned}a_{AA'A''} p^A q^{A'} &= 0 \\ a_{AA'A''} p^A r^{A''} &= 0 \\ a_{AA'A''} q^{A'} r^{A''} &= 0\end{aligned}\tag{1.3}$$

For our purposes, the important properties of the hyperdeterminant are that it is a quartic invariant under $[SL(2)]^3$ and under a triality that interchanges A, A' and A'' .

We recently found a physical application of this hyperdeterminant [2] by associating the eight components of $a_{AA'A''}$ with the four electric and four magnetic charges of the STU black hole in four-dimensional string theory [3] and showing that its entropy [4] is given by

$$S = \pi \sqrt{-\text{Det } a}.\tag{1.4}$$

The hyperdeterminant also makes its appearance in quantum information theory [5]. Let the three qubit system $AA'A''$ be in a pure state $|\Psi\rangle$, and let the components of $|\Psi\rangle$ in the standard basis be $a_{AA'A''}$:

$$|\Psi\rangle = \sum_{AA'A''} a_{AA'A''} |AA'A''\rangle\tag{1.5}$$

Then the three way entanglement of the three qubits A, A' and A'' is given by the *3-tangle* [6]

$$\tau_3 = 4|\text{Det } a|.\tag{1.6}$$

As far as we can tell [2], the appearance of the Cayley hyperdeterminant in these two different contexts of stringy black hole entropy (where the $a_{AA'A''}$ are integers and the symmetry is $[SL(2, Z)]^3$) and 3-qubit quantum entanglement (where the $a_{AA'A''}$ are complex numbers

and the symmetry is $[SL(2, C)^3]$ is a purely mathematical coincidence. Nevertheless, it has already provided fascinating new insights into the connections between strings, black holes, and quantum information [7].

In this paper, we provide a third physical application by allowing $a_{AA'A''}$ to represent the eight variables of the four-dimensional bosonic string $(\dot{X}^\mu, X'^\mu)$ and showing that in signature $(2, 2)$ the Nambu-Goto Lagrangian [8, 9] is given by $\sqrt{\text{Det } a}$. Interchange of the indices A , A' and A'' then reveals hitherto hidden discrete symmetries of the Nambu-Goto action.

2 The Nambu-Goto action

The Nambu-Goto action in a flat target space of signature $(2, 2)$ is given by

$$I_{NG} = -\frac{T}{2} \int d\tau d\sigma \sqrt{-\det \gamma} \quad (2.1)$$

where

$$\gamma_{A''B''} = \partial_{A''} X^\mu \partial_{B''} X^\nu \eta_{\mu\nu} \quad (2.2)$$

and $\eta = \text{diag } (1, 1, -1, -1)$. We introduce a two-component notation

$$X^{AA'} = \frac{1}{\sqrt{2}} \begin{pmatrix} -X^0 + X^2 & X^1 - X^3 \\ -X^1 - X^3 & -X^0 - X^2 \end{pmatrix} \quad (2.3)$$

and define

$$a_{AA'A''} = \begin{pmatrix} \dot{X}^{AA'} \\ X'^{AA'} \end{pmatrix}, \quad (2.4)$$

where $\dot{X} = \partial X / \partial \tau$ and $X' = \partial X / \partial \sigma$. Hence

$$\gamma_{A''B''} = \epsilon^{AB} \epsilon^{A'B'} a_{AA'A''} a_{BB'B''} \quad (2.5)$$

Then from (1.1) and (1.2)

$$\det \gamma = -\text{Det } a \quad (2.6)$$

The $SL(2, R)$ acting on the index A and the $SL(2, R)$ acting on the index A' are just the $O(2, 2) \sim SL(2, R) \times SL(2, R)$ spacetime symmetry, while the $SL(2, R)$ acting on the index A'' is the worldsheet symmetry

$$\begin{pmatrix} \dot{X}^\mu \\ X'^\mu \end{pmatrix} \rightarrow \begin{pmatrix} d & c \\ b & a \end{pmatrix} \begin{pmatrix} \dot{X}^\mu \\ X'^\mu \end{pmatrix} \quad (2.7)$$

where a, b, c, d are constants satisfying $ad - bc = 1$.

3 Discrete symmetries

Interchanging A and A' is just reverses the sign of $\dot{X}^1$ and X'^1 but the 4 transformations

$$\begin{aligned}
a_{AA'A''} &\rightarrow a_{A''AA'} \\
a_{AA'A''} &\rightarrow a_{A'A''A} \\
a_{AA'A''} &\rightarrow a_{A''A'A} \\
a_{AA'A''} &\rightarrow a_{AA'A'}
\end{aligned} \tag{3.1}$$

represent new discrete symmetries which are given explicitly by

$$\begin{pmatrix} \dot{X}^0 \\ \dot{X}^1 \\ \dot{X}^2 \\ \dot{X}^3 \\ X'^0 \\ X'^1 \\ X'^2 \\ X'^3 \end{pmatrix} \rightarrow \frac{1}{2} \begin{pmatrix} \dot{X}^0 - X'^1 - \dot{X}^2 + X'^3 \\ -X'^0 - \dot{X}^1 + X'^2 + \dot{X}^3 \\ -\dot{X}^0 - X'^1 + \dot{X}^2 + X'^3 \\ X'^0 - \dot{X}^1 - X'^2 + \dot{X}^3 \\ X'^0 + \dot{X}^1 + X'^2 + \dot{X}^3 \\ \dot{X}^0 - X'^1 + \dot{X}^2 - X'^3 \\ X'^0 - \dot{X}^1 + X'^2 - \dot{X}^3 \\ \dot{X}^0 + X'^1 + \dot{X}^2 + X'^3 \end{pmatrix} \tag{3.2}$$

$$\begin{pmatrix} \dot{X}^0 \\ \dot{X}^1 \\ \dot{X}^2 \\ \dot{X}^3 \\ X'^0 \\ X'^1 \\ X'^2 \\ X'^3 \end{pmatrix} \rightarrow \frac{1}{2} \begin{pmatrix} \dot{X}^0 + X'^1 - \dot{X}^2 + X'^3 \\ X'^0 - \dot{X}^1 - X'^2 - \dot{X}^3 \\ -\dot{X}^0 + X'^1 + \dot{X}^2 + X'^3 \\ X'^0 + \dot{X}^1 - X'^2 + \dot{X}^3 \\ X'^0 - \dot{X}^1 + X'^2 + \dot{X}^3 \\ -\dot{X}^0 - X'^1 - \dot{X}^2 + X'^3 \\ X'^0 + \dot{X}^1 + X'^2 - \dot{X}^3 \\ \dot{X}^0 - X'^1 + \dot{X}^2 + X'^3 \end{pmatrix} \tag{3.3}$$

$$\begin{pmatrix} \dot{X}^0 \\ \dot{X}^1 \\ \dot{X}^2 \\ \dot{X}^3 \\ X'^0 \\ X'^1 \\ X'^2 \\ X'^3 \end{pmatrix} \rightarrow \frac{1}{2} \begin{pmatrix} \dot{X}^0 - X'^1 - \dot{X}^2 + X'^3 \\ X'^0 + \dot{X}^1 - X'^2 - \dot{X}^3 \\ -\dot{X}^0 - X'^1 + \dot{X}^2 + X'^3 \\ X'^0 - \dot{X}^1 - X'^2 + \dot{X}^3 \\ X'^0 + \dot{X}^1 + X'^2 + \dot{X}^3 \\ -\dot{X}^0 + X'^1 - \dot{X}^2 + X'^3 \\ X'^0 - \dot{X}^1 + X'^2 - \dot{X}^3 \\ \dot{X}^0 + X'^1 + \dot{X}^2 + X'^3 \end{pmatrix} \tag{3.4}$$

$$\begin{pmatrix} \dot{X}^0 \\ \dot{X}^1 \\ \dot{X}^2 \\ \dot{X}^3 \\ X'^0 \\ X'^1 \\ X'^2 \\ X'^3 \end{pmatrix} \rightarrow \frac{1}{2} \begin{pmatrix} \dot{X}^0 + X'^1 - \dot{X}^2 + X'^3 \\ -X'^0 + \dot{X}^1 + X'^2 + \dot{X}^3 \\ -\dot{X}^0 + X'^1 + \dot{X}^2 + X'^3 \\ X'^0 + \dot{X}^1 - X'^2 + \dot{X}^3 \\ X'^0 - \dot{X}^1 + X'^2 + \dot{X}^3 \\ \dot{X}^0 + X'^1 + \dot{X}^2 - X'^3 \\ X'^0 + \dot{X}^1 + X'^2 - \dot{X}^3 \\ \dot{X}^0 - X'^1 + \dot{X}^2 + X'^3 \end{pmatrix} \tag{3.5}$$

4 Conclusions

One way to understand this triality is to think of the string as having three different world-sheets (Alice, Bob and Charlie) with three different metrics

$$\begin{aligned}\alpha_{AB} &= \epsilon^{A'B'} \epsilon^{A''B''} a_{AA'A''} a_{BB'B''} \\ \beta_{A'B'} &= \epsilon^{A''B''} \epsilon^{AB} a_{AA'A''} a_{BB'B''} \\ \gamma_{A''B''} &= \epsilon^{AB} \epsilon^{A'B'} a_{AA'A''} a_{BB'B''}\end{aligned}\tag{4.1}$$

All are equivalent, however, since

$$\det \alpha = \det \beta = \det \gamma = -\text{Det } a\tag{4.2}$$

It is remarkable that the Nambu-Goto action, first written down in 1970 can still reveal some secrets, although the $(2,2)$ signature means that their physical significance is not obvious. It is worth remarking, however, that this does form the bosonic sector of the $N=2$ critical superstring. See [10], for example.

Finally we recall [7] that the $E_{7(7)}$ Cartan invariant J_4 , depending on $28 + 28$ variables, reduces to Cayley's hyperdeterminant in a canonical basis. So one could generalize the Nambu-Goto action to

$$I_{NG} = -\frac{T}{2} \int d\tau d\sigma \sqrt{J_4}\tag{4.3}$$

Bearing in mind the embeddings

$$E_{7(7)} \supset SO(6,6) \times SL(2, R) \supset SO(2,2) \times SL(2, R)\tag{4.4}$$

this would describe a string in 28 dimensions with reduced Lorentz group $SO(6,6)$, rather than $SO(2,2)$. It may seem strange to embed the Lorentz group in a bigger group, but this was already true for the conventional Nambu-Goto theory (2.1), since $[SL(2, R)]^3$ and triality form a semi-direct product.

Perhaps our familiarity with string theory will allow us to gain yet more insights into the other two contexts in which Cayley's hyperdeterminant makes its appearance, namely black holes and quantum information, and vice-versa.

5 Acknowledgements

I am grateful to Dimitrios Giataganas for pointing out some typographical errors in the original version. I also thank Jussi Kalkkinen and Tom Kibble for the observation that $[SL(2, R)]^3$ and triality do not commute.

References

- [1] A. Cayley, "On the theory of linear transformations," Camb. Math. J. 4 193-209, 1845.

- [2] M. J. Duff, “String triality, black hole entropy and Cayley’s hyperdeterminant,” arXiv:hep-th/0601134.
- [3] M. J. Duff, J. T. Liu and J. Rahmfeld, “Four-dimensional string-string-string triality,” Nucl. Phys. B **459**, 125 (1996) [arXiv:hep-th/9508094].
- [4] K. Behrndt, R. Kallosh, J. Rahmfeld, M. Shmakova and W. K. Wong, “STU black holes and string triality,” Phys. Rev. D **54**, 6293 (1996) [arXiv:hep-th/9608059].
- [5] A. Miyake and M. Wadati, “Multipartite entanglement and hyperdeterminants,” [arXiv:quant-ph/02121146].
- [6] V. Coffman, J. Kundu and W. Wootters, “Distributed entanglement,” Phys. Rev. A **61** (2000) 52306, [arXiv:quant-ph/9907047].
- [7] R. Kallosh and A. Linde, “Strings, black holes, and quantum information,” arXiv:hep-th/0602061.
- [8] Y. Nambu, “Duality and hydrodynamics,” Lectures at the Copenhagen conference, 1970.
- [9] T. Goto, “Relativistic quantum mechanics of one-dimensional mechanical continuum and subsidiary condition of dual resonance,” Prog. Theor. Phys. **46** (1971) 1560.
- [10] H. Ooguri and C. Vafa “Geometry of $N = 2$ strings,” Nucl. Phys. B **361** (1991) 469.